Discuss issues associated with the classification and/or diagnosis of schizophrenia.

AO₁

DSM - IV

continuous disturbance for a least 6 months

ICD - 10

Doctors reluctant to diagnose schizophrenia

20% recover completely

10% significant improvement

30% some improvement

40% never recover

(10% commit suicide)

Population validity (cannot be generalised across cultures)

2.1% African Americans diagnosed 1.4% White Americans diagnosed

social norms or cultural biases can be an explanation

AO2

- P: 2 different diagnostic manuals, both have different requirements to diagnose schizophrenia
- E: Poor reliability of diagnosis and a la mof diagnosis a la mof diagnosis a la mof diagnosis a la mof diagnosis a
- Social stigma of schizophrenia and control of clark and schizophrenia and the other doesn't due to different factors measures to diagnose the condition.

 Schizophrenics are reluctant to see a doctor or diagnosis

- E: Estimated only 20% of schizophrenics have been diagnosed
- E: However, this could be culturally specific and so this assumption cannot be generalised to other cultures where the social stigma has a less dramatic effect
- L: More cross-cultural research is needed to determine the cultural validity of this findings.
- P: Different outcomes dependent on the individuals experience
- E: Low predictive validity (cannot predict the future outcome)
- E: This reduces the ability to be able to predict how the disorder will develop and the final outcome for the patients
- L: More research could be carried out to see if theres any correlation between certain schizophrenic characteristics and the future of the sufferer.

Golden Nuggets

DSM - IV & ICD - 10

Concurrent Validity

Social Stigma

Cultural Validity

Different Outcomes

Predictive Validity

Population Validity

Social Norms