

What is a range sketch?	19
1.6.2 Quadratic inequalities	20
What is the graphical method of solving quadratic inequalities?	20
What is the tabular method of solving quadratic inequalities?	20
1.7 Algebraic manipulation	20
1.7.1 Difference between two squares.....	20
What is the identity for this?	20
1.8 Graphs and graphical solutions to equations	20
1.8.1 Cubic functions	20
What is a cubic function?	20
What is the factorised form of this kind of function?	20
What shape is the graph of a cubic function?	20
1.8.2 Sketching cubic functions	21
What is the technique for sketching the graph of a cubic function?	21
1.8.3 Sketching $y = x^3$	21
What is the technique for sketching this graph?	21
When does this technique apply?	21
How does this differ if there are constants?	21
1.8.4 Reciprocal functions	21
What is a reciprocal function?	21
What are the two forms of reciprocal function?	21
What makes these forms different?	21
How are they different?	21
What is the technique for sketching the graph of this kind of function?	22
How is the graph of this type of curve different to the graphs of different types of curve? ...	22
What is the name given to the two lines that the two sections of the graph of a reciprocal function get progressively close to, but never touch?	22
1.8.5 Graphical solutions to equations	22
What is the technique for finding the points of intersection of two graphs?	22
1.9 Graph transformations.....	22
1.9.1 How is the graph of $f(x)$ transformed horizontally and vertically?	22
1.9.2 How is the graph of $f(x)$ scaled horizontally and vertically?	22
1.9.3 What is the technique for answering questions where you are asked to transform $y = f(x)$?	22
1.9.4 What is the technique for reflecting a graph of $f(x)$ in the axes?	23
2.2. Coordinate geometry in the (x, y) plane	23

2.1 Equation of a straight line	23
2.1.1 What are the identities for the equations for straight lines?	23
How are these identities related?	23
What is useful about the form $y = mx + c$?	23
2.1.2 What is the formula for calculating the gradient of a line given two points that lie on it?	23
2.1.3 What is the formula for finding the equation of a line given the gradient m and one of the points it passes through?	23
2.1.4 What is the formula for finding the equation for a straight line given two points that it passes through?	23
2.2 Conditions for parallel and perpendicular lines	23
2.2.1 What are the conditions for two lines to be parallel?	23
2.2.2 What are the conditions for one line to be perpendicular to another?	24
What is the mathematical way to check if two lines are perpendicular to each other?	24
Given the gradient of a line, how can the gradient of the line perpendicular to it be found?	24
3 Sequences and series	24
3.1 Sequences	24
3.1.1 Background	24
What is a sequence?	24
What is the name given to each number in a sequence?	24
Question Technique	24
When given part of a sequence of numbers, what is the method of finding more terms of the sequence?	24
3.1.2 The n th term of sequence	24
What is meant by the n th term of a sequence?	24
What is the formula for the n th term of a sequence?	24
How is this formula used?	24
3.1.3 Recurrence relationships	25
What is a recurrence relationship?	25
Why must the first term of a sequence be given when expressing a sequence as a recurrence relationship?	25
How is this different from expressing a sequence as an expression in n , where n is the number of the term?	25
Give an example of a sequence and its recurrence relationship	25
3.1.4 Arithmetic sequences	25
What is an arithmetic sequence?	25
How can an arithmetic series be shown using a recurrence relationship?	25
Question Technique	25

What is the third result?

To rationalise a fraction in the form $\frac{1}{a - (\sqrt{b})}$, multiply the numerator and the denominator by $a + (\sqrt{b})$

1.3 Quadratic functions

1.3.1 What is a quadratic function?

A quadratic function is a function that is in the form

$$y = ax^2 + bx + c, \text{ where } a \text{ is not equal to } 0.$$

This can also be written as

$$f(x) = ax^2 + bx + c$$

1.3.2 What are the characteristics of a graph of a quadratic function?

The graph is a curve.

The graph has only one turning point.

The graph is symmetrical on either side of the turning point

1.4 Quadratic equations

1.4.1 Meaning

What is a quadratic equation?

A quadratic equation is an equation in the form

$ax^2 + bx + c = 0$, where x is an unknown variable, and a , b and c are constants, known as the quadratic coefficient, the linear coefficient and the constant term respectively.

What are you doing when you are solving a quadratic equation?

You are finding the values of x for which the equation $ax^2 + bx + c = 0$ is true.

How many are there?

A quadratic equation has two solutions.

How can the solutions of quadratic equation be seen visually?

On the graph of a quadratic equation, the values of x when the curve intercepts the x -axis, that is, when $y = 0$, are the solutions to the quadratic equation.

Preview from Notesale.co.uk
Page 14 of 37

1.9.4 What is the technique for reflecting a graph of $f(x)$ in the axes?

A translation of $y = f(-x)$ is a reflection of the original curve in the y-axis

A translation of $y = -f(x)$ is a reflection of the original curve in the x-axis

2 2. Coordinate geometry in the (x, y) plane

2.1 Equation of a straight line

2.1.1 What are the identities for the equations for straight lines?

$y = mx + c$, where m and c are integers.

$ax + by + c = 0$, where a , b and c are integers.

How are these identities related?

They are interchangeable with each other. An equation in the form $y = mx + c$ can be transposed into the form $ax + by + c = 0$.

What is useful about the form $y = mx + c$?

The constant m is the gradient of the line, while c is the intercept of the line on the y-axis.

2.1.2 What is the formula for calculating the gradient of a line given two points that lie on it?

Given that the first point is at (x_1, y_1) and the second point is at (x_2, y_2)

$$m = (y_2 - y_1) / (x_2 - x_1)$$

2.1.3 What is the formula for finding the equation of a line given the gradient m and one of the points it passes through?

Given that the gradient is m and the point that it passes through is (x_1, y_1) ,

$$y - y_1 = m(x - x_1)$$

2.1.4 What is the formula for finding the equation for a straight line given two points that it passes through?

Given that the first point is at (x_1, y_1) and the second point (x_2, y_2) ,

$$(y - y_1) / (y_2 - y_1) = (x - x_1) / (x_2 - x_1)$$

2.2 Conditions for parallel and perpendicular lines

2.2.1 What are the conditions for two lines to be parallel?

Two lines are parallel if their gradients are the same.

where a_1 is the first term, and a_n is the last term in the series.

Why are there two forms for this equation?

The formula for the n th term of an arithmetic series is:

$$a_n = a_1 + (n-1)d$$

This value of the n th term can be substituted into the formula for the sum of an arithmetic series if the last term in the series is unknown, but the common difference is.

Similarly, if the common difference is unknown, but the last term is, you can use the simpler form of the equation.

Explain how this equation is derived.

Take the general arithmetic series as:

$$S = a + (a + d) + (a + 2d) + (a + 3d) + \dots + (a + (n-2)d) + (a + (n-1)d)$$

Reversing the sum:

$$S = (a + (n-1)d) + (a + (n-2)d) + (a + (n-3)d) + \dots + (a + d) + a$$

Adding the sums:

$$2S = [2a + (n-1)d] + [2a + (n-1)d] + [2a + (n-1)d] + \dots + [2a + (n-1)d]$$

There are n lots of $[2a + (n-1)d]$. That means we can express this sum as:

$$2S = n[2a + (n-1)d]$$

$$S = n/2 * [2a + (n-1)d]$$

3.2.4 Question Technique 2

What is the technique for answering a question where given an arithmetic series with its first term, common difference and length visible, you are asked to find the sum of the series?

1. Substitute the values into the formula for the sum of an arithmetic series
2. Simplify the equation to find the sum

What is the technique for answering a question where given an arithmetic series with its first term, last term and length, you are asked to find the sum of the series?

1. Substitute the value of n , a_1 and a_n into the second form of the equation - $S = n/2(a_1 + a_n)$
2. Simplify the equation to find the sum.

Is this the only thing that this gradient can represent? Explain your answer.

No. The straight line $y = 1/3x$ is not the only possibility. Any straight line with gradient $1/3$ may be written as $y = 1/3x + c$, where c is an arbitrary constant.

This is the most general equation which gives $dy/dx = 1/3$.

What does this mean that the equation $dy/dx = 1/3$ represents?

It means that equation represents all straight lines with a gradient of $1/3$.

What does it mean to find the anti-derivative with respect to x of this equation?

From the theorem, we know that antidifferentiation is equivalent to integration, so by taking the antiderivative with respect to x of the equation, we are finding the integral with respect to x . Thus, calculating the integral requires that we perform the reverse process of differentiation.

We know that the most general equation for y which when differentiated gives $dy/dx = 1/3$ is

$y = 1/3.x + C$, where C is an arbitrary constant.

Thus, for $dy/dx = 1/3$, the antiderivative, or integral is calculated as

$$\int dy/dx \, dx = \int 1/3 \, dx$$

Evaluating:

$$y = 1/3.x + C$$

5.1.3 What is the general formula for integration of x^n ?

$$\text{If } dy/dx = x^n$$

$$y = 1/n+1 * x^{n+1} + c$$

Why is this formula for integration?

As we now know, integration is the reverse process of differentiation. The formula for differentiation is:

$$\text{If } y = x^n, \, dy/dx = nx^{n-1}$$

The formula for integration has the function of providing the reverse process to this formula.

5.1.4 How is the formula for integration applied to a polynomial?

It is applied in the same way that the formula for differentiation applies to a polynomial. Integrate each term separately.

Preview from Notesale.co.uk
Page 36 of 37