## **STOICHIOMETRY AND PERCENT PURITY**

Many samples of chemicals are not pure. We can define percent purity as

 $\frac{\text{mass of pure compound in the impure sample}}{\text{total mass of impure sample}} \quad x \quad 100$ 

If an impure sample of a chemical of **known** percent purity is used in a chemical reaction, the percent purity has to be used in stoichiometric calculations. Conversely, the percent purity of an impure sample of a chemical of **unknown** percent purity can be determined by reaction with a pure compound as in an acid-base titration. Percent purity can also be determined, in theory, by measuring the amount of product obtained from a reaction. This latter approach, however, assumes a 100% yield of the product.

## Examples

Consider the reaction of magnesium hydroxide with phosphoric acid.

(b) Calculate the mass of 88.5%  $Mg(OH)_2$  needed to make 127 g of  $Mg_3(PO_4)_2$ , assuming a 100% yield.

mass Mg(OH)<sub>2</sub> = 127 g Mg<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub> x  $\frac{1 \text{ mole Mg}_3(PO_4)_2}{262.9 \text{ g Mg}_3(PO_4)_2}$  x  $\frac{3 \text{ moles Mg}(OH)_2}{1 \text{ mole Mg}_3(PO_4)_2}$  x  $\frac{58.3 \text{ g Mg}(OH)_2}{1 \text{ mole Mg}(OH)_2}$ = 84.49 g Mg(OH)<sub>2</sub>. mass 88.5% Mg(OH)<sub>2</sub> = 84.49 g Mg(OH)<sub>2</sub> x  $\frac{100 \text{ g 88.5\% Mg}(OH)_2}{88.5 \text{ g Mg}(OH)_2}$  = 95.5 g