


Staging by FIGO



Stage ①:-

- 1A1 < 3mm deep - TOC conization / family complete hysterectomy can be done
 ↳ type 1 (C)
 1A2 ≥ 3mm but < 5mm - Type 2 Hysterectomy
 (B) = modified Radical = w/out this also Pelvic lymph node - clony
 ↳ type 2 (C)
 ↳ Simple Excisional hyster
 ↳ X & N
 1B1 ≥ 5mm but < 2cm
 1B2 ≥ 2cm but < 4cm
 1B3 ≥ 4cm not Sx → chemo radiation

Stage ②:-

2A - invades upper 2/3 of vagina but not parametrium

A₁ < 4cm (Sx) → * Type 3 (C) = Radical hysterectomy + Pelvic lymphadenectomy
 A₂ ≥ 4cm (chemo radiation)

2B - invasion of parametrium
 all stages ≥ 2B - Chemoradiation

3A - spread to lower 1/3 of vagina

3B - spread to lat pelvic wall (Hydrocoelitis)

3C 1 the pelvic LN

2 + 10 paraaortic LN

4A - invasion bladder / rectum

on hysteroscopy → Bullous edema of mucosa → X 4A

B → distant spread

(inguinal node involved)

Preview from Notesale.co.uk
Page 6 of 7