## CHAPTER 2 : THE CELL:

CELL - The basic unit of living structure and function. The adult human body consists of more than 50 trillion cells. It is about 60% water, bathed by a solution called INTERSTITIAL FLUID.; They differ in shape and function.

## CELL PARTS:

- 1. NUCLEUS- located at the center of the cell. Control center of the cell. Contains the genetic material
- 2. CYTOPLASM- semi fluid that surrounds the nucleus. Site of cellular activities.
  - A. Cytosol semi- transparent fluid that suspends other elements, water is the main component.
  - B. Organelles metabolic machinery; specialized compartments each performing its own function to maintain the life of the cell.
  - > Ribosomes round dark bodies; sites of protein synthesis.
  - > Endoplasmic Reticulum (ER) fluid filled cisterns (tubes or canals); serves as a mini circulatory system for the cell.
    - 2 forms : Rough ER studded with ribososmes; export protein products Smooth ER - no ribosomes; for cholesterol breakdown, fat metabolism and detoxification of drugs.
      - Golgi Apparatus flat apparatus: sort proteins for transport for specific locations (mailing tag for proper destination)
      - > Lysosomes for intracellular digestion; present in WBC that engulf bacteria and harmful substance. (Pacman effect) 30
      - Peroxisomes sacs that detoxify poisonous substances (eg. Alcohol)
      - Mitochondria powerhouse of the cell. Energy.
      - Centrioles used in cell division.

Inclusions - substances that may or may not be present.

Eg. Melanin in skin / Melatonin

- 3. PLASMA MEMBRANE encloses the cell. A layer of proteins and phospholipids. > > (geni -per. neagle) specializations:
  - Microvilli fingerlike projections used for absorption.
  - > Junctions meeting point.
    - a. tight junctions impermeable; prevents passage of substances . (eg. Small intestines - prevent seeping into the bloodstream. )
    - b. desmosomes anchoring junctions (eg. Skin cells prevent con bulled apart that cause stress.)
  - c. gap junctions allow communication le Clear

1. DIFFUSION are mentifrom an area of higher to aver-concentration.
2. OSMOSIS Convement through a sm. Commende membrane.

Ca. hypertonic solution—absorb too much water. Swell.

- - - c. Isotonic solution equal. ( eg. Hospitalized isotonic sol. Replenish fluids in
- 3. BULK TRANSPORT too large to cross the membrane.
  - a. endocytosis folds into a pouch then encloses the particles
  - b. Pinocytosis cell drinking. Water droplets.
  - c. Phagocytosis Engulfs food ( solid materials).
  - d. Exocytosis opposite of endocytosis. Wastes leave the cell.

MITOSIS - cell division; should produce same material with the original cell.

## STAGES:

- 1. Prophase
- 2. Metaphase
- 3. Anaphase
- 4. Telophase

## CONTROL OF MITOSIS / TUMORS AND CANCER:

Causes of abnormal changes in cell:

- 1. heredity
- 2. certain chemicals
- 3. radiation

TUMOR - abnormal growths .

- a. benign cells stay together, removed by surgery.
- Malignant cells break free, migrate to new locations if touched.

CANCER - spread of malignant cells to new locations; abnormal size, shape and abilities (invade normal cells)

TREATMENT - chemotherapy or radiation therapy.