

### Case Presentation

It is the instance of an 18-year-old adolescent admitted to intensive outpatient services. The reason for doing this assessment is to collect early information about a patient in which the physician and the patient collaborate to establish what signs and symptoms are present and what they indicate. It had a significant impact on the later stages of treatment. The ASAM Criteria employs a multidimensional assessment method. This assessment is a tool to see how treatment might affect multiple living areas of a person. It has six dimensions that portray various aspects of the patient, and each represents a single aspect of the individual that contributes to the condition.

Dimension 1- Risk Rating: The patient reports withdrawal symptoms during his adolescence. The patient's last used substance was on September 21, 2021. He identified his withdrawal symptoms as 1/10. He noticed some trouble with sleeping and cold sweats. The patient reports severe intoxication and reports minimal withdrawal over the months.

Dimension 2 - Risk Rating: The patient reports slurred speech, eyesight problems, sleepiness, and a slow reaction time after consuming alcohol during withdrawal in a facility. The patient claims that he is consuming less food, resulting in supplement consumption.

Dimension 3 - Risk Rating: The patient reports that memory lapses are affecting his learning in academics, concentration, executive functions, recollection depth perception, and cognitive processing. The patient expressed that he tried to control his emotional suffering. The patient was cooperating with his sessions and participating in activities.

Dimension 4 - Risk Rating: The patient expresses a desire to break free from his addiction. He requires careful monitoring and getting a support system from the facility several times a week.