

72/ A patient with cystic fibrosis has a great amount of secretion at the lingula.

What is the ideal position that the therapist should put the patient in to drain this secretion?

- OA. Prone
- OB. Supine
- OC. Semi-lying
- OD. Side-lying on the right side

73/ A 58-year-old woman presented with severe hip pain. She has been taking steroids for her dermatomyositis. A CT scan hip confirmed avascular necrosis of femoral neck with an impacted fracture.

There is a 2 cm leg-length discrepancy.

Which of the following is the most appropriate intervention to reduce hip pain?

- OA. Shoe raise
- OB. Walking stick
- OC. Shortwave diathermy
- OD. Modified seating system

74/ A 67-year-old patient with lower back pain referred to the Physical Therapy Clinic for assessment and treatment. The therapist decided to start with the 1st performance of the pelvic tilt exercise.

Which of the following positions was performed?

- OA. Sitting
- OB. Standing
- OC. Lying supine on a firm surface
- OD. Lying supine on a soft surface

Preview from Notesale.co.uk
Page 9 of 15

75/ A 33-year-old patient with a reconstruction of massive tear of the rotator cuff. The Physical Therapist decided to initiate the full active shoulder flexion.

Which of the following is the most appropriate prescribed time in weeks to initiate the movement?

OA. 1

OB. 2

OC. 3

OD. 6

Preview from Notesale.co.uk
Page 10 of 15