

D- Acromioclavicular ligament doesn't contribute to stabilizers
Stabilizers are : Sternoclavicular joint and it's ligaments , interarticular disc

282-which part not included in posterior neck triangle

- a-clavicle
- b-trapezuis
- c-strenoclidomastoid
- d-strenum

The correct Answer is:
D- Sternum

ساعت فی الامدادات

د. احمد فتحي

د. محمد محمد

282-which is not related to the shoulder triangle

- a)Clavicle
- b)Sternum
- c) Humurars

The correct Answer is:
B- Sternum

283-Protraction and retraction of shoulder occurs at
a-glenohumeral joint
b-scapulothoracic interface
c-acromioclavicular joint

The correct Answer is:
B- Scapulothoracic interface

277-What is the action of this muscle shape ?



جروب معا لنتفوق في برومترك العلاج الـ

د. سيد صحصاد

د. محمد مرعي

- A- hip adduction
- B- hip extenstion
- C-back extension

The correct Answer is:

Preview from Notesale.co.uk
Page 7 of 20

YOUR BEST CHOICE

The correct Answer is:

B-Flexion isn't needed for spleneous capitis action which is extension , rotation , side bending

297-Splenius capitis response of :

- a- neck flexion
- b- neck extension
- c- neck rotation
- d- neck side bend

ساعت فی الامداد

د. احمد فتحي

د. محمد فرحات

The correct Answer is:

B-neck extension as we described above

298-which of following not needed in testing the trapezius muscle :

- a-head flexion
- b-head extension
- c-head rotation
- d-head lateral flexion

The correct Answer is:

A-head flexion isn't needed in testing the trapezius because it's action is rotation, retraction, elevation, and depression of scapula

299-which of following not needed in testing the sternocleidomastoid muscle:

- a-head flexion
- b-head extension
- c-head rotation
- d-head lateral flexion

The correct Answer is:

A- head extension , sternocleidomastoid action Unilaterally; cervical rotation to opposite side, cervical lateral flexion to same side Bilaterally; cervical flexion, raises the sternum and assists in forced inhalation

د. سيد صحتاح

د. محمد فرحات

300-A physical therapist is beginning an evaluation of a 5-year-old boy. The mother indicates that she pulled the child from a seated position by grasping the wrists. The child then experienced immediate pain at the right elbow. The physician's orders are for right elbow range of motion and strengthening. Which of the following is the most likely diagnosis?

312-A patient with a diagnosis of a rotator cuff tear has just begun active range of motion. The therapist is strengthening the rotator cuff muscles to increase joint stability and oppose the superior shear of the deltoid. Which of the rotator cuff muscles participate least in opposing the superior shear force of the deltoid?

- a- Infraspinatus**
- b- Subscapularis**
- c-Teres minor**
- d-Supraspinatus**

The correct Answer is:
D- supraspinatus

313-A therapist is attempting to gain external rotation range of motion in a patient's right shoulder. The therapist decides to use contract-relax-contract antagonist. In what order should the following rotator cuff muscles contract to perform this movement successfully?

- a-Infraspinatus - teres minor**
- b-Subscapularis – supraspinatus**
- c-Teres minor – infraspinatus**
- d-Supraspinatus – subscapularis**

The correct Answer is:
D- supraspinatus – subscapularis

314--in condromalacia consider in the ttt?

- a- patellar tapping and strenthening of the quadriceps**
- b- strenthening vastus medialis only**
- c- strenthening vastus lateralis only**
- d- strenthening of hamstring**

The correct Answer is:
A- patellar Tapping and strentheninh of the quadriceps