

B- Chondrosarcoma

348-most common site of estrogenic sarcoma is:

- a-Femur, upper end**
- b-Femur, lower end**
- c-Tibia, upper end**
- d-Tibia, lower end**

**The correct Answer is:
A- Femur, Upper end.**

349-Deformity associated with colles fracture (fracture lower end of radius)

- a- Dinner fork deformity**
- b - Pes planus**
- c- Talipes equinovarus**

**The correct Answer is:
A- Dinner fork deformity**

350- Colles fracture may possibly cause injury to which of the following

- a- flexor pollicis**
- b- abductor pollicis**
- c- adductor pollicis**
- d- extensor pollicis longus**

**The correct Answer is:
D- Extensor pollicis longus.**

351-What the other name of fracture radius:

- a-no name**
- b-pontine fracture**
- c-bennt fracture**
- d-colles fracture**

**The correct Answer is:
D- Colles fracture**

352-which is incorrect about plump line of body in standing

The correct Answer is:

B- 120 Degree as we described above

366-The angle between longitudinal axis passing through femoral neck and longitudinal axis of femur is decreased in the case of:

A -genu valgum

B -genu varum

C -genu recurvatum

D -coxa Vera

ساعت فی الامداد

د. احمد فتحي

د. محمد صبرى

The correct Answer is:

D- Coxa Vera as we described

367-Observing a patient in a standing position, the therapist notes that an angulation deformity of the right knee causes it to be located medially in relation to the left hip and foot. This condition is commonly referred to as:

a-genu varum

b-genu valgum

c-pes cavus

d-none of the above

The correct Answer is:

B- genu Valgum

**Preview from Notesale.co.uk
Page 10 of 20**

368-the normal angle between the femur and the neck of femur is 126 when the angle increase the deformity is:

a- coxa valga

b- genu Valgus

C- genu recurvatum

The correct Answer is:

A- Coxa Valga as we described above.

جروب معا لنتفوق فى برومترك العلاج الطبيعى

369-Coxa Vera present as abnormality of

a-proximal femur

b-distal femur

c-proximal tibia

d-distal tibia

د. سيد صبرى

د. محمد مرسى

The correct Answer is:

A- proximal Femur

The correct Answer is:

B- Lateral trunk flexion & convex side to RT

391-A 14-year-old girl with right thoracic scoliosis is referred to physical therapy. The therapist should expect which of the following findings?

- a-Left shoulder high, left scapula prominent, and right hip high**
- b-Left shoulder low, right scapula prominent, and left hip high**
- c-Right shoulder high, right scapula prominent, and right hip high**
- d-Right shoulder low, right scapula prominent, and left hip high**

The correct Answer is:

B- left shoulder lo

392-The most common deformity in geriatric is

- a- Kyphosis**
- b- Scoliosis**
- c- Kyphoscoliosis**
- d- Lordosis**

The correct Answer is:

A- Kyphosis

393- Severe kypho-scoliosis will lead to

- a- Left rather than right ventricular failure**
- b- Right rather than left ventricular failure**
- c- Frequent respiratory infections**
- d- No cardiac abnormalities**

The correct Answer is:

C- frequent respiratory infections

394-You are evaluating an athlete who is complaining of pain in the left shoulder region. Your assessment of the shoulder elicits pain in the last 30 degrees of shoulder abduction range of motion. This finding is most congruent with

- a- calcific supraspinatus tendinitis.**
- b- sub acromial bursitis.**
- c- acromio-clavicular sprain.**
- d- thoracic outlet syndrome.**

The correct Answer is:

C- Acromio-Clavicular sprain due it`s range of motion