

Row-Reduced Echelon (REE) Form

Examples: Solve the following system using Gauss-Elimination Method

Preview from Notesale.co.uk Page 6 of 40



Eigenvalues of $4 4^{-1} - 4 2^{-1}$	4
$Eigenvalues of 4A = 4\lambda$	1 2
_	=,-2
	5 <i>′</i>
	5
Eigenvalues of $4^2 - 1^2$	25 4
Eigenvalues of $A = \lambda$	25,4
-	
Examples of $A^2 = 2A + I = 12 = 21 + 1$	1(0
Eigenvalues of $A^ ZA + I = \lambda^ Z\lambda + I$	10,9
Eigenvalues of $43 + 2I = 13 + 2$	107 (
Eigenvalues of $A^2 + ZI = \lambda^2 + Z$	12/,-0
Ŭ	

Example 2:Find the eigen values of $A = \begin{bmatrix} 3 & 2 \\ 3 & 8 \end{bmatrix}$

Solution: Given $A = \begin{vmatrix} 3 & 2 \\ 3 & 8 \end{vmatrix}$, then the characteristic equation of matrix A is



-2 -12^{-1} -8 **Example 3**: Find the eigen values and eigen vector of the matrix A= 1 4 4 1 J 0 Solution: 8/3 4 0 1 0 1 0 0 = The characteristic equation is $|A - \lambda I_n| = 0$ $0 \mid 0 \rfloor R_1 \rightarrow R_1 - 8/3R_2$ 0 0

Let *A* be $n \times n$ matrix and λ be an eigen value for *A*. If λ occurs $(k \ge 1)$ times then *k* is called the **Algebraic multiplicity** of λ , and the number of basis vectors is called **Geometric multiplicity**.

Example: Find eigen values and eigen vectors of the matrix. $A = \begin{bmatrix} -2 & 2 & -3 \\ 2 & 1 & -6 \\ -1 & -2 & 0 \end{bmatrix}$. Also determine algebraic and geometric multiplicity. <u>Solution</u>: The characteristic equation is $|A - \lambda I_n| = 0$. $\begin{vmatrix} 2 & \lambda & 2 & -5 \\ 2 & 1-\lambda & -6 \\ -1 & -2 & 0-\lambda \end{vmatrix}$ $= (-2 - \lambda)[(1 - \lambda)(-\lambda) - (-2)(-6)] - 2[2(-\lambda) - (-1)(-6)] - 3[2(-2) - (-1)(1 - \lambda)]$ $= (-2 - \lambda)[-\lambda + \lambda^{2} - 2] - 2[-2\lambda - 6] - 3[-4 + 1 - \lambda]$ $= -\lambda^{3} - \lambda^{2} + 21\lambda + 45$ $= -(\lambda^{3} + \lambda^{2} - 21\lambda - 45)$ $\therefore -(\lambda^{3} + \lambda^{2} - 21\lambda - 45) = 0$ $\therefore (\lambda + 3)(\lambda^{2} - 2\lambda - 15) = 0$ $\Rightarrow (\lambda + 3)(\lambda + 3)(\lambda - 5) = 0$ $\Rightarrow \lambda_1 = 5, \lambda_2 = -3, \lambda_3 = -3$ Algebraic Multiplicity of $\lambda = -3$ is 2 and of $\lambda = 5$ is 1. We solve the following homogeneous system: $\therefore [A - \lambda I]X = \begin{vmatrix} -2 - \lambda & 2 & -3 \\ 2 & 1 - \lambda & -6 \\ -1 & -2 & 0 - \lambda \end{vmatrix} \begin{vmatrix} x_1 \\ x_2 \\ x_3 \end{vmatrix} = \begin{vmatrix} 0 \\ 0 \\ 0 \end{vmatrix}$

Case I: When $\lambda_1 = 5$	Case II : When $\lambda_2 = -3$, $\lambda_3 = -3$

Theorem:Every square matrix can be decomposed as a sum of symmetric and skew-symmetric matrices.

Proof: Let *A* be $m \times n$ matrix.

Let $B = \frac{1}{2}(A + A^T)$ and $C = \frac{1}{2}(A - A^T)$ be two matrices.

Obviously, A = B + C

Now,
$$B^T = \left[\frac{1}{2}(A + A^T)\right]^T = \frac{1}{2}[(A + A^T)]^T = \frac{1}{2}[A^T + (A^T)^T] = \frac{1}{2}(A^T + A) = B$$

As $B^T = B$, B is symmetric.

$$C^{T} = \left[\frac{1}{2}(A - A^{T})\right]^{T} = \frac{1}{2}[(A - A^{T})]^{T} = \frac{1}{2}[A^{T} - (A^{T})^{T}] = \frac{1}{2}(A^{T} - A) = -C$$

Therefore, $C^T = -C$, C is skew-symmetric.

Therefore, A is a sum of symmetric and skew-symmetric matrices.

Calcy –Hamilton Theorem Every square matrix satisfies its own characteristic equation **A** for the orem states that, for a square matrix *A* of order *n*, if $|A - \lambda I_n| = 0$. **Note that Example (i)**: Verify Calcy Hendrilton theorem and cerife and the inverse of $A = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}$ and A^4 . Solution: the characteristic equation for given matrix is $|A - \lambda I_2| = 0$. $\Rightarrow \begin{vmatrix} 1 - \lambda & 4 \\ 2 & 3 - \lambda \end{vmatrix} = 0$ $\Rightarrow (1 - \lambda)(3 - \lambda) - 8 = 0$ $\Rightarrow \lambda^2 - 4\lambda - 5 = 0$ Now, by putting $\lambda = A$, we have $A^2 - 4A - 5I = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix} - 4 \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix} - 5 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 9 & 16 \\ 8 & 17 \end{bmatrix} - \begin{bmatrix} 4 & 16 \\ 8 & 12 \end{bmatrix} - \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0$

Hence, Cayley-Hamilton theorem verified.

Now, by using Cayley-Hamilton theorem, we have